

Patient Name:	Date of Birth:
Date of Appointment:	Appointment Time:
Referring Physician:	
Appoir	ntment Location:
MAIN OFFICE: 1222 S. Patterson Blvd.—Suite 400 Dayton, OH 45402 (937) 222-0022	■ ENGLEWOOD OFFICE: 9000 N. Main Street—Suite 319 Dayton, OH 45415 (937) 832-3582
CENTERVILLE OFFICE: 6601 Centerville Business Pkwy.—Suite 200 Dayton, OH 45459 (937) 435-7476	☐ <b>TROY OFFICE:</b> 1861 Towne Park Drive—Suite H Troy, OH 45373 (937) 222-0022
-	st Information Sheet:
	nine the cause of your symptoms. You will be having the following tests:
of the middle and inner ear.  WHAT TO EXPECT: You will be placed in a sound	deproof booth with a set of headphones over the ears or small
	asked to convey that you have heard the tone by either st may also conduct speech tests whereby you will be asked to repeat the
-	
· · · · · · · · · · · · · · · · · · ·	
□ <u>VEMP(Vestibular Evoked Myogenic Potential</u>	): (Testing Code: 92700)
<b>PURPOSE:</b> To measure the function of your balance	e nerve. gh on your sternum (chest) and on your neck. You will be asked to turn you
□ VNG or ENG (Video- or Electro-nystagmogra	aphy): (Testing Codes: 92540 & 92537)
<b>PURPOSE:</b> To test the inner ear's balance function. <b>WHAT TO EXPECT:</b> A recording will be made of y	your eye movements in different conditions. During this test we will ask you ons, and we will also bathe each ear canal with warm and cool air. You may

experience some brief periods of dizziness. Rest assured, the dizziness will subside quickly and in except for rare cases you should

have no difficulty driving home after the test. *TIME*: This test will last about 1 hour.



## Special Instructions for <u>VNG or ENG</u> Testing:

Certain substances can influence the body's response to testing, causing an invalid response.

Therefore, please follow the following test instructions.

DAY OF TESTING:

No food or drink four (4) hours before the test (including caffeinated products) to avoid an upset stomach during testing.

Please do not wear eye makeup, foundation, or moisturizer on your face.

Dress comfortably, dresses and skirts should be avoided. Slacks and shoes without heels should be worn.

48 HOURS PRIOR TO TESTING:

No alcohol 48 hours prior to testing. This includes any quantity of beer, wine, liquor, and cough medicines containing alcohol.

The following medications may affect the results of the VNG or ENG: If stopping the medication is approved by the prescribing physician, the medications should be not be taken for 48 hours prior to the VNG. They can be resumed immediately after the testing. If these medications cannot be discontinued, under a physician's direction, the testing can still be performed.

- Anti-Vertigo Medication: Antivert, Dro-Peridol Drops, Meclizine, Ru-vert, etc.
- Anti Nausea Medication: Atarax, Bontrol, Bonine, Bucladin, Compazine, Dramamine, Marezine, Phenergan, Thorazine, Scopalomine Patch, etc.
- <u>Tranquilizers:</u> Atarax, Atarazz, Ativan, Elavil, Equanil Etraafon, Klonipin, Librium, Librax, Lorazapam, Miltown, Prozac, Serax, Travil, Tranxene, Valium, Vistaril, Xanax, etc.
- Muscle Relaxers: Flexeril, etc.
- <u>Sedatives:</u> Butisol, Chloral Hydrate, Dalmane, Doridan, Halcion, Nembbutal, Restoril, Placidly, "pm" pain medications, Quaalude, Seconal, Sleeping pills, etc.
- Narcotics & Barbituates: Codeine, Darvocet, Darvon, Demerol, Dilaudid, Florinal, Oxycontin, Oxycodone, Percocet, Percodan, Phenaphen, Rebaxin, Tylenol with Codeine, Vicodin.
- Anti-Histamines: Actifed, Allegra, Benadryl, Chlor-Trimeton, Claritin, Comtrex, Dimetane, Dimetapp, Disophrol, Drixoral, Polaraminee, Seldane, Tavist, Teldrin, Temaril, Triaminic, Trinalin, Vistaril, Zyrtec, and any over-the-counter cold remedies, etc.
- Anti-Seizure Medications: Dilantin, Tegretol, Phenobarbital, etc.

This is not a complete list of such medications, so please ask if there are questions about any medications being taken

If any of these medications have been taken by the patient for psychiatric, seizure, or other neurologic disorders, they should only be discontinued after consulting with the prescribing physician as stopping these medications suddenly may result in withdrawal symptoms or other serious medical conditions.

You should continue to take any medications prescribed for you except those categories listed above. You should continue to take all heart and blood pressure medications, hormone medication, Thyroid Medications, Diabetes or Sugar Medications. You may take vitamins, Aspirin, Bufferin, Tylenol, etc.

IF ANY QUESTIONS ARISE REGARDING TEST PROCEDURES OR MEDICATIONS  $\underline{PLEASE\ CALL\ THE\ OFFICE}$ 

(PLEASE COMPLETE THE QUESTIONNAIRE ON THE OPPOSITE PAGE)



Name:	
Date of Birth:	

## **DIZZINESS/VERTIGO QUESTIONNAIRE**

wim	headed ming Sensation	Objects spinning or turning around you	Loss of Balance Confusion			
Tendency to fall to the Right Tendency to fall to the Left		Spinning inside you Feeling like you will pass out	OTHER:			
	· ·	cribed (above) do you experience any of	f the following? (Check all that			
ply).		Г. В.	F1 1'			
Naus		Ear Pain	Flushing Numbness			
Vomi Head	•	Visual Changes Spots before your eyes	Loss of Body Control			
	le Vision	Heart pounding in your chest	Nervousness			
	ng or Noises in your ears.	Difficulty Speaking	Muscle Weakness			
_	ng Loss	Difficulty Swallowing	Falling Suddenly			
	ressure	Hunger	Loss of Consciousness			
ease	describe your symptoms furthe	r•				
1)	• • •	-time occurrence? Recurrent attack	κs?			
2)	How many occurrences of syn	nptoms have you had?				
3)	How long do your symptoms	last?				
4)	When was the first occurrence	?				
5)						
6)	Have you had any history of e	ar infections, ear surgery, or ear problen	ns?			
7)						
8)						
9)						
10	) Do your symptoms get worse	before or after eating?				
11]	) Do your symptoms get worse	when you are tired or stressed?				
12	) Have you had any treatment for	or dizziness?				
		dizziness, with what?				
14	) Did this treatment for dizzines	s help?				
		of dizziness?				
17	) Have you ever been exposed t	o any irritating fumes?				
		rmation which you feel is important that				
		, 1				